



YaraAmplix™
OPTITRAC™

NOTICE

Before using this product, read the entire label

OPTITRAC™

(7-0-2)

Liquid micro-nutrient with extract of *Ascophyllum nodosum* for foliar application to agricultural crops.

LIQUID FERTILIZER

Guaranteed analysis:

7% w/v (70 g/L) Total nitrogen (N)

[6% w/v as urea 1% w/v as organic]

2.2% w/v (22 g/L) Total potassium (K) as hydroxide

1.3% w/v (13 g/l) Boron (B) as ethanalamine salt

1.3% w/v (13 g/L) Zinc (Zn) as sulphate

Impurities:

Biuret 18.7g biuret/kg N

!! Warning Statements !!

Do not swallow. The spray from this product may act as an irritant.

Avoid inhalation and contact with the eyes and skin.

Users are encouraged to seek further advice.

This fertilizer contains biuret.

Repeated foliar sprays may cause damage to citrus.

10 litres
Net weight
11.7 kg

Distributed by **Yara Australia Pty. Ltd.**, Level 2, 200 Mary Street, Brisbane QLD 4000, Australia

Tel: 0299594266

www.yara.com.au

Manufactured by **Yara UK Ltd.**, Pocklington Industrial Estate, Pocklington, York, YO42 1DN, UK.

YaraAmplix™ OPTITRAC™

RECOMMENDED APPLICATION RATES AND TIMINGS

Almond: Apply 2 - 3 l/ha when the first leaves appear, at the start of flowering and at the end of flowering / fruit swelling. Water rate: 200 l/ha minimum.

Apple, Apricot, Cherry, Nectarines, Peach, Pears, Plum: 2 - 3 l/ha applied pre-flowering, at flowering and also post-petal fall. Water rate: 200 l/ha minimum.

Avocado: 2 - 3 l/ha before flowering, at flowering, and at fruit set. Water rate: 200 l/ha minimum.

Bananas: 3 l/ha applied to new leaf growth in the spring and repeated twice at 10 to 14 day intervals. Water rate: 30 to 500 l/ha. Alternatively: 6 l/ha per year in total. Apply 12 sprays of 0.5 l/ha per year. Do not spray on developing fruit.

Beans: 2 - 3 l/ha from the 2 to 3 leaf stage up to the flower buds stage (pre-flowering). Water rate: 200 l/ha minimum.

Blueberries: 2 - 3 l/ha. Apply at the beginning of flowering, repeating at full flowering and fruit set. Also, apply during periods of stress, repeating at 10 to 14 day intervals as necessary. Water rate: 200 l/ha.

Brassicas, Soya bean: Three applications of 2 - 3 l/ha at 10 to 14 day intervals commencing at the 4 to 6 leaf stage. Water rate: 200 l/ha minimum.

Carrot: Three applications of 3 l/ha at 10 to 14 day intervals commencing when the crop is 15 cm tall. Water rate: 200 l/ha minimum.

Cereals: 1-3 l/ha applied from the 2 leaf stage to first node detectable (Zadoks G.S. 12 to 31), with repeat applications within these stages at 10 to 14 day intervals, as necessary. Water rate: 200 l/ha minimum.

Citrus: 3 l/ha applied pre-flowering, at flowering and during fruit formation. Water rate: 200 l/ha minimum.

Cotton: 2-3 l/ha applied at flower bud squares and again at open flowers. Water rate: 200 l/ha minimum.

Cucurbits (Field grown), Pepper / Capsicum and Tomato (Field grown), Onion: Three applications of 3 l/ha at 10 to 14 day intervals commencing at the 4 to 6 leaf stage. Water rate: 200 l/ha.

Grapes (table), Grapes (wine): 3 l/ha applied pre-flowering, at flowering and also at fruit set. Water rate: 200 l/ha minimum. Alternatively: 3 l/ha at flower truss visible, at flower buds separated / start of flowering and at fruit set. Water rate: 200 l/ha minimum.

Hazelnuts, Walnuts: Apply 3 l/ha when the first leaves appear, at the start of flowering and at the end of flowering / fruit swelling. Water rate: 200 l/ha minimum.

Leafy vegetable: Max Rate of Single Application is 3 l/ha; Water rate: 200 l/ha; 2 weeks after transplanting or for direct sown crops at the 4 to 6 leaf stage. Repeat as necessary at 7 to 14 day intervals.

Oilseed rape / canola: 1 - 3 l/ha at the 4-9 leaf stage and again at the onset of the rapid stem extension stage up to the start of flowering. Water rate: 50-200 l/ha.

Olive: 3 l/ha at bud break, 10 to 14 days post-flowering and during fruit formation. Water rate: 200 l/ha minimum.

Potatoes: 3 l/ha applied 7 to 14 days after 100 percent emergence, 20 days after tuber initiation and, following petiole analysis, during tuber bulking. Water rate: 200 l/ha minimum.

Protected crops: 0.3 litres per 100 litres water maximum concentration. Water rate: 1000 l/ha maximum. Refer to equivalent field grown crop for application timing.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable clothing, gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the SDS for further information, available at www.yara.au

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.


CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



5 033704 019271 >

YaraAmplix™ OPTITRAC™



UN 3082

ENVIRONMENTALLY HAZARDOUS
SUBSTANCE, LIQUID, N.O.S. (Zinc sulphate)

DANGER

Hazardous ingredients: glycerol, zinc sulphate (monohydrate), 2-aminoethanol, 2-methylisothiazol-3(2H)-one

H318: Causes serious eye damage. H315: Causes skin irritation. H317: May cause an allergic skin reaction. H411: Toxic to aquatic life with long lasting effects. P280: Wear protective gloves/clothing and eye/face protection.

P261: Avoid breathing gas, vapor or spray. P305+P351+P338+P310: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician. P302+P352+P333+ P313: IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs, seek medical advice/attention. P391: Collect spillage.

Yara Australia Pty. Ltd., Level 2, 200 Mary Street, Brisbane QLD 4000, Australia
Tel: 0299594266 Emergency telephone number (24/7) +61 2801 44558